

Nightguard Home Care – Patient Instructions

“What you can do for yourself may be more important than what we can do for you.”

Control Muscle Abuse

- Avoid sleeping on your jaw
 - ☒ Try a cervical pillow
- Use your appliance as directed
 - ☐ Every time you sleep
 - When not wearing it, clean it, then store in the case
- Avoid chewy foods or any foods that fatigue your muscle
 - ☐ Ice, raw vegetables, hard crusted bread, apples, and large sandwiches
- Avoid clenching your teeth
 - ☐ “Lips together, teeth apart”
- Consider muscle strengthening exercises

Type of Stretching	Main Characteristics	Parameters	Purpose
Static stretching (passive)	A specific position is held with the muscle in tension to a point of a stretching sensation	10-30 s (<90 s), 2-4 repetitions	Improve range of motion prevent injury improve performance
Active dynamic stretching	Gently propelling the muscle toward the maximum range of motion	3-6 repetitions of 15-30 s	Improve viscoelasticity improve range of motion

- Do not open too wide
- Limit caffeine within 6 hours of bedtime
 - ☐ Healthy diet, adequate sleep, and exercise
- Apply moist heat/massage twice a day
- Take medication as prescribed
 - ☐ Ibuprofen
 - ☐ Naproxen
 - ☐ Other

KOIS CENTER, LLC
1001 FAIRVIEW AVE N.
SUITE 2200
SEATTLE, WA 98109

TEL 206.621.5310
FAX 206.621.7609
WEB KOISCENTER.COM